

HOW DO WE EVALUATE THE VULNERABILITIES OF MIGRATORY SPECIES TO CLIMATE CHANGE?

Hector Galbraith
National Wildlife Federation



VULNERABILITY ASSESSMENT

- ◆ Some species may be relatively easy




OTHERS PRESENT MAJOR DIFFICULTIES - SUPERMIGRANTS



RED KNOT MIGRATION AND STOPOVER SITES



MIGRATORY WILDLIFE VULNERABILITY ASSESSMENT

- ◆ Difficult challenges for VA:
 - Highly extravagant lifestyles
 - Where? Breeding range, wintering range, stopover sites, migration itself, all of above?
 - Synchronicity?
 - Data hard to come by from parts of range
 - Wind?
- 
- A stylized, dark teal silhouette of a mountain range is positioned in the bottom right corner of the slide, partially overlapping the bottom edge of the text area.

RED KNOT – WHERE ARE THE VULNERABILITIES?

- ◆ Tierra del Fuego?
- ◆ Argentina coast?
- ◆ Brazil?
- ◆ Mid-Atlantic states?
- ◆ Hudson's Bay?
- ◆ High Arctic?
- ◆ Fall or spring?
- ◆ Wind patterns?
- ◆ Synchronicities?

Comprehensive VA Needed?

A stylized, dark teal mountain range graphic is located in the bottom right corner of the slide, extending from the right edge towards the center.

Vulnerabilities of Shorebirds to Climate Change

Hector Galbraith¹, Stephen Brown²,
David DesRochers³, Michael Reed⁴

¹National Wildlife Federation

²Manomet Center for Conservation Sciences

³Dalton State College

⁴Tufts University


A stylized, dark teal mountain range graphic is located in the bottom right corner of the slide, extending from the right edge towards the center.

Objectives


- ◆ Evaluate change in extinction risk of North American shorebirds due to climate change



Why Shorebirds?

- ◆ Reported widespread declines
 - ◆ Sentinels of global environmental change because of their hemispheric ecosystem use
 - ◆ Migratory aggregations of some species are a spectacular biological phenomenon
 - ◆ Iconic species valued by public?
- 
- A stylized, dark teal silhouette of a mountain range is positioned in the bottom right corner of the slide, adding a decorative element to the background.

SHOREBIRDS ARE IN TROUBLE



Based on migration counts in eastern N.America; Bart et al 2007. *J Av. Biol*

MAIN QUESTION ASKED

- ◆ How much does climate change move the needle on the existing vulnerability categories of USCP/PIF?

U.S. Shorebird Conservation Plan

Risk Categories

- 1) Not at Risk
- 2) Low Concern
- 3) Moderate Concern
- 4) High Concern
- 5) Highly Imperiled
- 6) ~~Holy Smokes! Really,~~
~~highly imperiled~~
Critical



Vulnerability Factors

	Score	Arrow
1) Loss/gain in breeding habitat under climate change	3	↑
2) Loss/gain in wintering habitat under climate change	5	↑↑
3) Loss/gain in migration habitat under climate change	3	↑
4) Degree of dependence on ecological synchronicities	5	↑↑
5) Migration distance	4	↑
6) Degree of breeding, wintering, or migration habitat specialization	4	↑↑

Risk Factors

1) Loss/gain in breeding habitat under climate change:

	Score	Arrow
Major loss (>50%)	5	↑↑
Moderate loss (10-50%)	3	↑
Limited or no loss (-10-10%)	0	0
Moderate increase (10-50%)	-1	↓
Major increase (>50%)	-2	↓↓

Note: risk could decrease

Example: Semipalmated Sandpiper



1) Loss/gain in breeding habitat under climate change:

	Score	Arrow
Moderate loss (10-50%)	3	↑

Yearlong rainfall predicted to ↑ throughout breeding range. May result in flooding & loss of much breeding habitat especially since the species prefers drier areas with access to water. Nesting habitats along shorelines also could ↓ as a result of increased rainfall.

Confidence = low




Semipalmated Sandpiper

	Score	Arrow
1) Loss/gain in breeding habitat under climate change	3	↑
2) Loss/gain in wintering habitat under climate change	5	↑↑
3) Loss/gain in migration habitat under climate change	3	↑
4) Degree of dependence on ecological synchronicities	5	↑↑
5) Migration distance	4	↑
6) Degree of breeding, wintering, or migration habitat specialization	4	↑↑

Change in status from 'moderate concern' to 'highly imperiled'

Application

- ◆ Evaluated 49 species of shorebird breeding in North American north of Mexico
 - ◆ For each factor, included confidence level
 - ◆ Determined shifts in risk categories
- 
- A stylized, dark teal silhouette of a mountain range is positioned in the bottom right corner of the slide, partially overlapping the text of the third bullet point.

Results for 50 Shorebirds

- ◆ 43 species (86%) predicted to ↑ risk level due to climate change
 - 34 increased by 1 level
 - 9 increased by 2 levels
- ◆ 3 species at lower risk
 - ◆ Solitary sandpiper – more breeding habitat
 - ◆ Bristle-thighed curlew – more breeding & wintering habitat
 - ◆ White-rumped sandpiper – more wintering habitat



U.S. Shorebird Conservation Plan

Risk Category	Current	Expected with climate change
Not at risk	0	0
Low concern	7	2
Moderate concern	15	7
High concern	23	13
Highly imperiled	4	17
Critical	–	10

Species in New 'Critical' Category

- ◆ Snowy Plover
- ◆ Wilson's Plover
- ◆ Piping Plover
- ◆ Mountain Plover
- ◆ Am. Oystercatcher
- ◆ Long-billed curlew
- ◆ Bar-tailed godwit
- ◆ Ruddy turnstone
- ◆ Sanderling
- ◆ Short-billed dowitcher




Where from here?



- ◆ Detailed species-specific VA
- ◆ ID common risks as focus for management activity
 - e.g., shoreline habitat on migration routes & wintering areas
- ◆ Still reviewing the assessments & considering degree of threat to shift risk category
- ◆ We welcome feedback, things to consider, insights, information

TAKE HOME MESSAGES

- ◆ For complex spp. We need complex, comprehensive VA
 - ◆ They are doable (?)
 - ◆ Build off of existing structures if possible (PIF, NAWP, etc.)
 - ◆ Must be resilient to lack of data
 - ◆ Can they be applied to less well-known species?
- 
- A stylized, dark teal silhouette of a mountain range is positioned in the bottom right corner of the slide, partially overlapping the bottom edge of the text area.